

A NEW METHOD FOR REARING *LUTZOMYIA TRAPIDOI* (DIPTERA: PSYCHODIDAE), WITH OBSERVATIONS ON ITS DEVELOPMENT AND BEHAVIOR IN THE LABORATORY¹

By **Byron N. Chaniotis²**

Abstract: A new technique for rearing the Panamanian phlebotomine sand fly *Lutzomyia trapidoi* (Fairchild & Hertig) is described. Experimental data are provided on sugar- and blood-feeding, mating, oviposition and culture of immature stages. As many as 70% of females survived oviposition when maintained in styrofoam chambers and fed concentrated sucrose solutions.

¹This work was partially supported by NIH Contract NIAID 72-2514.

²Gorgas Memorial Institute of Tropical and Preventive Medicine, Middle America Research Unit, Box 2011, Balboa Heights, Canal Zone.

Lutzomyia trapidoi (Fairchild & Hertig) is widely distributed in Panama (Fairchild & Hertig 1952). It bites a wide variety of vertebrates, including man (Chaniotis et al. 1971, Tesh et al. 1972), and is known to be a vector of cutaneous leishmaniasis (Johnson et al. 1963), as well as a number of arboviruses (Tesh et al. 1974). Attempts to colonize this species have not been successful. Hertig & Johnson (1961) and Johnson & Hertig (1961) in a major effort to rear Panamanian sand flies succeeded

