

## EXOERYTHROCYTIC STAGES OF *PLASMODIUM VIVAX* IN *ATELES* MONKEYS\*

DAVID C. BAERG, RICHARD N. ROSSAN, AND MARTIN D. YOUNG

Gorgas Memorial Laboratory, Apartado 6991, Panama 5, Republic of Panama†

**Abstract.** Seven and 10-day exoerythrocytic forms of *Plasmodium vivax* have been experimentally produced in hepatic tissues of *Ateles* (spider monkeys); patent infections developed in the recipients. These monkeys represent a new test system.

In 1969, for the first time, exoerythrocytic schizonts of a human malaria (*Plasmodium falciparum*) were demonstrated in a New World monkey (*Aotus trivirgatus*).<sup>1</sup> Later, in two brief reports,<sup>2,3</sup> it was stated that these forms of *P. vivax* also had been seen in experimentally infected *Aotus* monkeys. A detailed description of 6-, 7-, and 8-day bodies in the vivax-*Aotus* model has since been published.<sup>4</sup>

Subsequent to the transmission of vivax malaria to *Ateles fusciceps* (black spider monkeys),<sup>5</sup> we have found 7- and 10-day exoerythrocytic stages of the parasite in this second host system. The present communication will record our preliminary observations.

### MATERIALS AND METHODS

Sporozoites of the Achiote line of *P. vivax* were derived from *Anopheles albimanus* after infective feedings on *Aotus* donors. Mosquito thoraces were triturated in chilled 10% homologous, heat inactivated serum saline (0.85%) and sporozoites obtained by centrifugation. Three *A. fusciceps* served as recipients; each had been splenectomized and was free of naturally acquired malaria. Hepatic inoculation of sporozoites and biopsies on each monkey were performed via laparotomy. The liver tissue was fixed in Carnoy's solution, and processed by routine histological procedures. Specimens were embedded in

paraffin, and serially sectioned at 5  $\mu$ . Colophonium-Giemsa was used for staining.<sup>6</sup>

### RESULTS AND DISCUSSION

Exoerythrocytic stages were recorded for each animal (Table 1). In the biopsy material from 328D, a single 7-day form was identified, whereas for 488D, 9 schizonts have been observed. In the latter case, examination of the tissue is continuing. No 10-day forms were found in these two monkeys, though patent infections did appear 2 and 12 days, respectively, after the second biopsy. In the third subject, 5833, two exoerythrocytic forms were seen on day 10, the first day of parasite patency. The second biopsy from this monkey on day 15, was negative for parasites.

The 7-day bodies were circular or ovoid in appearance, ranging in maximum diameter from 15.0  $\mu$  to 25.0  $\mu$  (mean = 20.1  $\mu$ ). The number of nuclei at this stage of development was variable, 4 to 30 per section. The nuclei were deeply stained, and in most cases were of uniform size, discrete and rounded. The cytoplasm was slightly granular, compact, with several small, clear vacuoles in approximately half of the parasites. Shrinkage from the host cell was noted in a few instances. Compression of the host cell nucleus occurred, and there was no apparent tissue reaction in the surrounding area.

The 10-day exoerythrocytic stages were lobed, with longest diameters measuring 63.8  $\mu$  and 67.5  $\mu$ . A section of an unlobed portion of one of these forms is shown in Figure 1. Nuclei were finely divided, irregularly-shaped and not clearly defined. The cytoplasm was extremely granular and diffuse; beginning cleft formation was apparent. One of the bodies was more heavily stained on one side. Small vacuoles also were present in one body. There was moderate shrink-

Accepted 5 January 1974.

\* This work was sponsored by the U.S. Army Medical Research and Development Command under Grant No. DADA 17-72C-2031. This is contribution number 1236 to the Army Research Program on Malaria.

† Mailing address: Gorgas Memorial Laboratory, P. O. Box 2016, Balboa Heights, Canal Zone.

