

THE EPIDEMIOLOGY OF HEPATITIS B ANTIGEN AND ANTIBODY AMONG PANAMANIAN CUNA INDIANS*

W. C. REEVES,† C. J. PETERS,‡ AND R. H. PURCELL

Gorgas Memorial Institute of Tropical and Preventive Medicine, Middle America Research Unit, Balboa Heights, Canal Zone, and National Institute of Allergy and Infectious Diseases, Laboratory of Infectious Diseases, National Institutes of Health, Bethesda, Maryland 20014

Abstract. Previous studies of hepatitis B antigen (HBsAg) and antibody to it (anti-HBs) showed widely differing exposures between Panamanian Indian tribes. Cuna Indians living on islands appeared infrequently exposed to HBsAg; we found no one antigenemic and low age specific anti-HBs rates. In contrast, mainland dwelling Guaymi and Chocó Indians had a high prevalence of anti-HBs. We have now measured HBsAg by counter-electrophoresis and anti-HBs by radioimmunoassay in two Cuna Indian groups who live in the Darien forest. The prevalence of HBsAg among Darien Cuna was low, 3 positive of 239 tested, but 106 (44%) had anti-HBs. Darien Cuna thus evidenced greater exposure to HBsAg than island Cuna (8% had anti-HBs) and had an anti-HBs prevalence similar to the neighboring Chocó Indians (42%). The Guaymi Indians of western Panama had a lower frequency of anti-HBs (29%) than either Chocó or mainland Cuna but their frequency of chronic antigenemia was significantly greater. These data suggest that while exposure may be a function of village habitat, chronic antigenemia may reflect differences in host responses.

Exposure to hepatitis B antigen (HBsAg) varies considerably among Panamanian Indians. In a previous study, mainland dwelling Guaymi and Chocó Indians had a high frequency of antibody to HBsAg (anti-HBs) (26 and 42%) whereas only 8% of island Cuna Indians were positive.¹ It was difficult to account for these differing exposures to HBsAg, since the Indian groups did not practice self-mutilation nor were injections common. Village habitat constituted the most striking disparity between the three groups. The Guaymi live in western Panama's tropical deciduous forest and savannah; Chocó Indians live in the Darien rain forest; the previously sampled Cuna Indians occupy coral islands off eastern Panama's Caribbean coast. However, the Cuna Indian tribe also has villages close to the Chocó in the Darien forest. The present paper describes

the seroepidemiology of HBsAg and anti-HBs among the forest dwelling Cuna.

METHODS

Blood samples were obtained from the inhabitants of two clusters of mainland Cuna villages. In 1967 Drs. Pedro Galindo and Carl Johnson (Gorgas Memorial Laboratory) collected 99 sera during a Chagas disease survey in the Chucunaque River villages of Nura, Huala and Mortí. During 1972 we sampled 140 inhabitants of Maje, Icanti, Ipeti and Ibesipgana on the Bayano River. In each village we established medical clinics to vaccinate children against measles and provide medical care, and everyone 6 months or older attending the clinic had blood specimens taken. We also attempted to bleed all relatives of those attending the clinic.

Sera were assayed for HBsAg using counter-electrophoresis² and a guinea pig antiserum.³ An aliquot from each serum was tested for anti-HBs at the Laboratory of Infectious Diseases, NIAID, using the radioimmunoprecipitation technique.^{4,5}

Statistical analysis of HBsAg and anti-HBs frequencies utilized the Mantel Haenszel method,⁶ which combines individual 2 × 2 tables for each age group and controls them according to relative sample size and percent positive.

Accepted 7 February 1975.

* Supported by NIH-NIAID contract 72-2514.

Address reprint requests to: W. C. Reeves, Department of Epidemiology, SC-36, University of Washington, Seattle, Washington 98195.

† Present address: Departments of Medicine and Epidemiology-International Health, University of Washington, Seattle, Washington 98195.

‡ Present address: Scripps Clinic and Research Foundation, La Jolla, California 92037.

