

ON THE VALIDITY OF *HAEMAGOGUS SPEGAZZINII* FALCO KUMM
ET AL, 1946

(Diptera, Culicidae)

PEDRO GALINDO

Gorgas Memorial Laboratory, Apartado 1252, Panamá, R. de P.

Kumm *et al.* (1946) described a new *Haemagogus* from Colombia with hairy larva and male with short palps and bushy antennae, which could be clearly differentiated from *H. capricornii* Lutz, but appeared very close to *H. spegazzinii* Brethes, being separable from the latter species only by details of the mesosome of the male when viewed from the side. The new form was named *falco*, but the describers considered it doubtful whether the characters defining it were of sufficient importance to justify the creation of a new species and placed it as a subspecies of *spegazzinii*. In the same publication the name *H. janthinomys* Dyar, 1921, based on a species from the island of Trinidad, was relegated to the synonymy of *H. spegazzinii spegazzinii*.

Levi Castillo (1956), after examining the male terminalia of the holotype of *H. janthinomys*, concluded that this specimen shows the mesosome tip as in *falco*, and consequently sank *H. spegazzinii falco* Kumm *et al.*, 1946, in the synonymy of *H. janthinomys* Dyar, 1921.

The author is not in agreement with Levi Castillo's conclusions for the following reasons:

1) Cerqueira (1943) published a photomicrograph of the male terminalia of the holotype (then cotype) of *H. janthinomys*. As can be observed in this excellent reproduction, the terminalia is only partly dissected and the mesosome is in ventral view, thus not well oriented to judge the shape of its tip, which, as pointed out by Kumm *et al.* (*loc. cit.*), can best be studied in lateral view.

2) The author, on a visit to the U. S. National Museum, personally examined the holotype slide of the terminalia of *janthinomys*. Although the mesosome in this preparation is not in a good position to permit a conclusive judgment, it appears to belong to an intergrading form between *spegazzinii* and *falco*, close to the so-called "intermediate" type illustrated by Kumm and Cerqueira (1951) from areas of intergradation in Brazil.

3) Kumm *et al.* (*loc. cit.*) in the same paper in which they described *H. spegazzinii falco* from Colombia stated: "the name *janthinomys* becomes a synonym of *H. spegazzinii*, as material obtained from the type area of *spegazzinii*, near Ledesma, Argentina, is the same as that from the island of Trinidad, B.W.I., the type locality of *janthinomys*." This statement shows rather convincingly that these authors had material from Trinidad, as well as from Argentina, in front of them at the time they described *falco*, and found that specimens from Trinidad (*janthinomys*) were closer to type material of *spegazzinii* than to their new form.

4) The terminalia of 10 specimens of "*janthinomys*" from Trini-

dad in the collection of the Gorgas Memorial Laboratory have been dissected by the author and the mesosome mounted in lateral view. These specimens all show some degree of intergradation between *spgazzinii* and *falco*, but appear to be much closer to the former than to the latter (see photomicrographs).

5) More than a hundred mesosomes of males from Honduras, Nicaragua, Costa Rica and Panama examined by the author appear identical with material from Colombia (*falco*) and differ from the mesosomes of males from Trinidad (*janthinomys*.)

6) In mapping out the distribution of *spgazzinii* and *falco*, Kumm and Cerqueira (loc. cit.) show that while *spgazzinii* is quite abundant along the northeastern coast of Brazil, being found all the way up into the State of Amapá along the border with French Guiana, *falco* is not coastal at all but ranges throughout the northwestern corner of Brazil. Specimens from the island of Trinidad would be logically expected to fall closer to *spgazzinii*, the common form along the Atlantic littoral of northern Brazil, rather than to *falco* which is more Andean in distribution.

From these observations the author concludes that typical *falco* extends from northwestern Brazil and adjoining territories in Ecuador and Perú, through Western Venezuela and Colombia as far north as the north coast of Honduras. True *spgazzinii* occurs from northern Argentina and Bolivia, across eastern Brazil to French Guiana. The area of north-central Brazil, most of the Guianas, part of Venezuela, and the island of Trinidad (type locality of *janthinomys*) form a large intergrading zone where mesosomes of intermediate type are found, which neither correspond to typical *spgazzinii* nor to typical *falco*. Since the form *janthinomys* falls in this category, but appears closer to the former than to the latter, the author feels that there is no justification for sinking *H. spgazzinii falco* Kumm et al., 1946, in the synonymy of *H. janthinomys* Dyar, 1921, and suggests that the latter be maintained in the synonymy of *H. spgazzinii spgazzinii* Brethes, 1912, as proposed by Kumm et al. (loc. cit.), and that *falco* be considered a valid name to designate a northern and western geographical race of *spgazzinii*.

EXPLANATION OF PLATE

Fig. A, *H. spgazzinii spgazzinii*. Mesosome of a male from "Rio de Janeiro, Brazil (without date) J. Lane"; fig. B, *H. janthinomys*. Mesosome of a male from "St. Pat's, Arima, Trinidad, (8-15)-12-54. T. H. G. Aitken." (Note similarity with mesosome shown in figure A); fig. C, *H. janthinomys*. Mesosome of a male from Tabaguite, Trinidad, (8-13)-1-55. T. H. G. Aitken"; fig. D, *H. spgazzinii falco*. Mesosome of a male from "Choreha, Chiriqui, Panama, 5-6-50. P. Galindo." (Note differences with figures B and C); fig. E, *H. spgazzinii falco*. Mesosome of a male from "Rio Mesapa, El Negrito, Department of Yoro, Honduras, 4-9-54. P. Orjuela. (Northernmost specimen of the species thus far collected).

**B****C****A****D****E**

REFERENCES

- Cerqueira, N. L., 1943. Algumas espécies novas da Bolívia, e referencia a tres espécies de *Haemagogus*. Mem. Inst. Oswaldo Cruz 39: 1-14.
- Kumm, H. W., and Cerqueira, N. L. 1951. The *Haemagogus* mosquitoes of Brazil. Bul. Ent. Res. 42: 169-181.
- Kumm, H. W., Osorno-Mesa, E., and Boshell-Manrique, J. 1946. Studies on mosquitoes of the genus *Haemagogus* in Colombia. Amer. Jour. Hyg. 43: 13-28.
- Levi-Castillo, R. 1956. A systematic note on *Haemagogus' spegazzinii* Brethes, 1912. Proc. Ent. Soc. Wash. 58: 345-347.