

## TRYPANOSOMA CRUZI: EFFECTS OF RHODNIUS PROLIXUS EXTRACTS ON IN VITRO DEVELOPMENT

Ducll E. WOOD (\*) and Octávio E. SOUSA (†)

### SUMMARY

The addition of saline extracts of adult *Rhodnius prolixus* bugs to an insect-oriented culture system for *Trypanosoma cruzi* resulted in the large scale evolution of trypomastigote forms of these parasites *in vitro*. Parasite populations in these cultures reached densities of as many as  $8.3 \times 10^7$  parasites per ml after 14 days incubation at 27-28°C.

### INTRODUCTION

For many years research on the metabolism and immunology of *Trypanosoma cruzi* was limited because the available *in vitro* culture systems produced mainly epimastigote stages. In 1967 CASTELLANI et al.<sup>1</sup> reported the *in vitro* evolution of metacyclic trypomastigote forms after a series of manipulations through two types of media. WOOD & PIPKIN<sup>2</sup> subsequently reported that by using a system based on the insect cell culture medium of GRACE<sup>2</sup> (GMA), large numbers of trypomastigote stages of *T. cruzi* could be harvested. However, the insect-oriented culture required the addition of 0.5% of a specially processed hemolymph from diapausing pupae of the moth *Philosamia cynthia*. This requirement limited the availability of the insect-oriented culture system to only a few laboratories.

A search for a more available source of the insect factor(s) supplied in the moth hemolymph led to the following study.

### METHODS AND MATERIALS

A myotropic strain of *T. cruzi* (Brazilian origin) was used in these experiments. The strain was maintained in Swiss albino mice by serial blood passage every 10 days.

Adult *Rhodnius prolixus*, reduviid vectors of *T. cruzi*, were obtained from a laboratory colony, and extracts (Re) were prepared in the following manner. Bugs were ground in a constantly-chilled (4°C) mortar and pestle at a ratio of one adult per ml of phosphate buffered saline, pH 6.8 (PBS). After thorough grinding, the triturate was filtered through PBS soaked, Whatman's #50 filter-paper, in a Buchner funnel. The filtered extract was collected in a chilled Erlenmeyer flask, and then heat-inactivated at 56° C for 6 minutes. The material was then transferred into centrifuge tubes and frozen at -70° C for 12 hours. Following this, the extract was quick-thawed and centrifuged at  $14,000 \times G$  for 30 minutes at 4° C. The supernatant was sterilized by filtration through a 0.22 $\mu$  membrane filter and stored at -20° C.

(1) This investigation was supported in part by the Bureau of Medicine and Surgery and the Office of Naval Research Contract N.º N00014-73-C-0108

(2) The opinions or assertions contained herein are those of the Authors and are not to be construed as reflecting the views of the Navy Department or the Naval Service at large

Gorgas Memorial Laboratory

Apartado 6991 — Balboa Heights, Canal Zone

Panamá, Republic of Panamá

